

**Original Article**

## Prolonged apnea/hypopnea during water swallowing in patients with amyotrophic lateral sclerosis

Sonoko Nozaki, M.D.<sup>1)\*</sup>, Shuhei Sugishita, SLP<sup>2)</sup>, Toshio Saito, M.D.<sup>3)</sup>,  
Yoshifumi Umaki, M.D.<sup>1)</sup>, Katsuhito Adachi, M.D.<sup>1)</sup> and Susumu Shinno, M.D.<sup>3)</sup>

**Abstract:** Purpose Swallowing difficulty is increased along with progression of respiratory disturbance in patients with Amyotrophic Lateral Sclerosis (ALS). To analyze the respiratory patterns during swallowing is important for the management of this disease. In this study, we evaluated apnea/hypopnea during water swallowing and the respiratory cycle at rest and after water swallowing.

Method We evaluated respiratory patterns in swallowing in 10 ALS patients (66.0 ± 7.1 years old), in 10 Myotonic dystrophy (MD) patients (46.5 ± 12.2 years old), and in 10 healthy volunteers as control subjects (61.7 ± 10.0 years old). The ALS and MD patients had consulted the Department of Neurology of Toneyama National Hospital or Tokushima National Hospital between April 2002 and July 2006. Respiratory patterns were evaluated by simultaneous recording of cervical swallowing sound in water swallow. A hypersensitive microphone measured cervical sound. A thermister was used for pneumography. The means of four continuous respiratory cycles at rest and after swallow of 3 ml water were used for analysis. Respiration with amplitude of 1/2 or smaller than that of the pneumography at rest was defined as hypopnea, and the apnea/hypopnea duration was evaluated as the respiratory suppression time.

Statistical Analysis All analyses were performed using SPSS 11.0J (SPSS Inc., Chicago, IL).

Results In the ALS group, the respiratory cycle was 3.15 ± 0.76 sec (2.31-4.39 sec) at rest, while after swallowing, it was 2.78 ± 0.83 sec (1.77-4.80 sec) (p = 0.1). In the MD group, the respiratory cycle was 2.56 ± 0.46 sec (1.91-3.67 sec) at rest, while after swallowing, it was 2.94 ± 0.60 sec (2.03-4.29 sec). In the control group, it was 3.46 ± 0.57 sec (3.18-4.34 sec) at rest and 3.24 ± 0.50 sec (2.64-4.04 sec) after swallowing. The apnea/hypopnea duration during water swallow was 14.33 ± 8.89 sec (2.50-30.68 sec) in the ALS group, 3.66 ± 1.58 sec (1.78-6.42 sec) in the MD group, and 3.64 ± 1.00 sec (2.34-5.56 sec) in the control group. The apnea/hypopnea duration in the ALS group was significantly longer than that in MD and control groups (p = 0.005, p = 0.004 by the t-test). The ALS patients with severe respiratory failure or with aspiration in videofluoroscopy showed extended apnea/hypopnea duration.

Conclusion Prolonged apnea/hypopnea was observed during water swallowing in ALS patients. We speculate that this prolongation is caused by severe swallowing disturbance and respiratory failure, which increases the risk of aspiration. The respiration of ALS patients should be closely monitored during eating.

(臨床神経, 48 : 634—639, 2008)

**Key words :** amyotrophic lateral sclerosis, myotonic dystrophy, dysphagia, apnea, hypopnea

### Introduction

Amyotrophic lateral sclerosis (ALS) is an extremely se-

vere and progressive neuromuscular disease. Respiratory insufficiency and dysphagia develop as the disease progresses. Swallowing difficulty is increased along with progression of respiratory disturbance in patients with ALS. Therefore,

\*Corresponding author: School of Rehabilitation, Hyogo University of Health Science [1—3—6 Minatojima, Chuo-ku, Kobe, Hyogo, 650-8530 Japan]

<sup>1)</sup>Department of Neurology, National Hospital Organization, Tokushima National Hospital

<sup>2)</sup>Department of Rehabilitation, National Hospital Organization, Tokushima National Hospital

<sup>3)</sup>Department of Neurology, National Hospital Organization, Toneyama National Hospital

(Received: 27 December 2007)




Fig. 1 The cervical swallowing sound and the respiratory cycle simultaneously recording system.

analysis of respiratory patterns in swallowing is important for the management of this disease.

In this study, we evaluated apnea/hypopnea during water swallowing and the respiratory cycle before and after swallow in patients with ALS, in patients with myotonic dystrophy (MD), and in healthy volunteers (controls). Cervical sounds during swallowing were recorded while pneumography was simultaneously performed.

## Methods

We evaluated respiratory patterns during swallowing in 10 ALS patients, in 10 MD patients, and in 10 healthy volunteers. The ALS patients ( $66.0 \pm 7.1$  years old [range 57-77]), MD patients ( $46.5 \pm 12.2$  years old [range 26-60]), and control subjects ( $61.7 \pm 10.0$  years old [range 52-79]) had consulted the Department of Neurology of Toneyama National Hospital or Tokushima National Hospital between April 2002 and July 2006.

ALS was diagnosed according to the EL Escorial criteria<sup>1)</sup> and MD was diagnosed clinically. The control subjects had no cerebrovascular or neuromuscular disease, and did not exhibit choking (suggesting dysphagia), sputum sticking, or moist voice. No abnormalities were observed in this group during testing. Of the 10 ALS patients, 2 required supplemental tube feeding (ALS Functional Rating Scale swallowing part<sup>2)</sup>: FRSsw 0), 5 required a diet of better consistency (FRSsw 2), 2 occasionally choked (FRSsw 3), and 1 was able to swallow and eat normally (FRSsw 4).

The ethics committees of both institutions approved this study, and written consent was obtained from all subjects.

### Measurements

Respiratory patterns in water swallow were evaluated by simultaneous recording of cervical swallowing sound and

pneumography. A sensor, commonly used for measurement of sleep apnea, was used for pneumography because this is an established method for measuring respiratory curves without causing discomfort. Cervical sound was measured by a hypersensitive microphone.

Subjects sat quietly without talking or moving for 5 minutes, while resting respiration was continuously recorded. The operator marked events such as dry swallows, coughing, and body movements, which were deleted from the respiratory trace before analysis. The mean of four continuous respiratory cycles at rest was used for analysis. The same process was used to measure the respiratory cycle after the subjects drank 3 mL of water 3 times at 20-second intervals. Because dry swallows or coughs sometimes occurred, a sequence of four respiratory cycles with the least noise obtained during the 3 water swallows was used for the analysis. Again, the mean of four continuous respiratory cycles after swallowing was used for analysis.

A thermistor, TR-711T (L type) by Nihon Koden, was used for pneumography in this study, but the temporal changes from exhalation to inhalation and apnea were slightly unclear. Therefore, respiration with an amplitude of 1/2 or smaller than that of the pneumography at rest was defined as hypopnea, and the apnea/hypopnea duration was evaluated as the respiratory suppression time.

Swallowing and Respiratory Monitoring Equipment (Fig. 1)

Cervical sound<sup>3)</sup> was measured by a hypersensitive microphone (Microphone UC-92H by Sound level meter Unit UN-04 RION).

For the pneumography recording, a thermistor (5-mm diameter) was placed near, but not in contact with, the nostril (to avoid body temperature changes). The thermistor was connected to a respiratory/plethymo coupler (DAT Data re-

**Table 1** Profiles of the ALS patients, the MD patients, and the control subjects

	age (yo)	gender	%FVC (%)	VF measure
ALS Case 1	57	M	53	oral phase disturbance
ALS Case 2	59	F	46	unremarkable
ALS Case 3	59	M	81	unremarkable
ALS Case 4	59	F	88	unremarkable
ALS Case 5	66	M	24	penetration
ALS Case 6	68	F	130	unremarkable
ALS Case 7	69	M	80	aspiration
ALS Case 8	73	M	35	aspiration
ALS Case 9	73	F	34	aspiration
ALS Case 10	77	M	NIV	penetration
MD Case1	27	M	82	poor lingual movement
MD Case2	30	M	47	poor lingual movement
MD Case3	33	M	61	nd
MD Case4	44	F	43	aspiration
MD Case5	50	M	26	sever aspiration
MD Case6	54	F	74	nd
MD Case7	54	F	38	poor lingual movement
MD Case8	56	M	32	poor lingual movement, residue on pharynx
MD Case9	57	M	60	nd
MD Case10	60	M	52	poor lingual movement, aspiration
				water swallowing test (screening test)
Control 1	52	F	nd	normal
Control 2	53	F	nd	normal
Control 3	53	F	nd	normal
Control 4	55	F	nd	normal
Control 5	56	F	nd	normal
Control 6	58	F	nd	normal
Control 7	66	F	nd	normal
Control 8	70	M	nd	normal
Control 9	75	M	nd	normal
Control 10	79	M	nd	normal

FVC: forced vital capacity, VF: videofluoroscopy, nd: not done

corder RD-135T TEAC), and the cervical swallowing sound and the respiratory cycle were recorded simultaneously. Analysis was performed using the Multipurpose Bioelectric Signal Analysis Program BIMUTAS<sup>®</sup>II (Kissei Comtec Co., Ltd.).

Forced vital capacity was measured before the study.

#### Statistical Analysis

All analyses were performed using SPSS 11.0J (SPSS Inc., Chicago, IL). The t-test was used to compare the two data sets. The individual mean respiratory cycle of four respiratory cycles was used for analysis.

## Results

Profiles of the ALS patients, the MD patients, and the control subjects are shown in Table 1. Fig. 2 shows the results of the swallowing sounds and pneumography in a control subject, an ALS patient and a MD patients.

The respiratory cycle is expressed as mean  $\pm$  standard deviation (minimal value-maximal value) for each group. In the ALS group, the respiratory cycle was  $3.15 \pm 0.76$  sec (2.31-4.39 sec) at rest, while after swallowing, it was  $2.78 \pm 0.83$  sec (1.77-4.80 sec) ( $p = 0.1$ ). In the MD group, the respiratory cycle was  $2.56 \pm 0.46$  sec (1.91-3.67 sec) at rest, while after swallowing, it was  $2.94 \pm 0.60$  sec (2.03-4.29 sec). In the control group, it was  $3.46 \pm 0.57$  sec (3.18-4.34 sec) at rest and  $3.24 \pm 0.50$  sec (2.64-4.04 sec) after swallowing (Fig. 3).

The apnea/hypopnea during water swallow was  $14.33 \pm 8.89$  sec (2.50-30.68 sec) in the ALS group,  $3.66 \pm 1.58$  sec (1.78-6.42 sec) in the MD group, and  $3.64 \pm 1.00$  sec (2.34-5.56 sec) in the control group. The apnea/hypopnea duration in the ALS was significantly longer than that in MD and in control ( $p = 0.005$ ,  $p = 0.004$  by the t-test) (Fig. 4).

Some of the ALS and the MD patients with exhibiting respiratory failure show the tendency of the extended apnea/hypopnea (Fig. 5). There was no significant relationship be-




Fig. 2 The results of the swallowing sound and pneumography in an ALS patient, in a MD patient and in a control subject.




Fig. 3 The respiratory cycle before and after water swallowing (Awa) in ALS, MD and control. In the ALS group, the respiratory cycle was decreased after swallowing ( $p = 0.1$ ).

tween apnea/hypopnea and respiratory cycle after water swallow (Fig. 6).

Results of videofluoroscopy, which was done within 2 weeks, showed longer apnea/hypopnea times in ALS patients who had aspiration than in those without aspiration ( $p = 0.5$ ) (Fig. 7).

### Discussion

In this study, we evaluated apnea/hypopnea during water






Fig. 4 The apnea/hypopnea (A/H) duration in the ALS was significantly longer than that in MD groups and in the control group ( $p = 0.005$ ,  $p = 0.004$  by the t-test).

swallowing in patients with ALS, in patients with MD, and in control subjects; prolonged apnea/hypopnea was observed in ALS patients. In Hiest's study, the respiratory cycle after swallowing was decreased in healthy elderly subjects in an upright position<sup>4</sup>. In our study, we found the trend of the reduction of post swallow respiratory cycle in ALS ( $p = 0.1$ ), but could not find any significant change of respiratory cycle before and after water swallow in MD and control. Preiksatis compared one respiratory cycle immediately before swallow and one immediately after swallow, and showed the respiratory cycle in post swallow was slightly longer than that in pre swallow in younger subjects<sup>5</sup>. In supine position, Nishino<sup>6</sup> found that the respiratory cycle did not change af-



**Fig. 5** Relationship between apnea/hypopnea (A/H) duration and % forced vital capacity (%FVC) in ALS and MD patients. The apnea/hypopnea (A/H) duration was extended in some of the patients exhibiting respiratory failure.




**Fig. 6** Apnea/hypopnea (A/H) & respiratory cycle after water swallow (Asw) in ALS & MD.

ter water swallow in comparison with that before swallow in normal subjects. The differences and disagreements described in post swallow respiratory cycle durations may be due to methodological factors such as volume of water, position, analysis duration of respiratory cycle.

We showed extremely prolonged apnea/hypopnea in ALS patients during water swallowing. Three mechanisms for apnea/hypopnea prolongation during water swallowing in the ALS patients were considered:

1) Respiration was depressed by aspiration or penetration



**Fig. 7** Apnea/hypopnea and aspiration in videofluoroscopy (VF): ALS patients who had aspiration in VF showed longer apnea/hypopnea times than those without aspiration ( $p = 0.5$ ). MD patients did not show that relationship.

during the swallow. Results of videofluoroscopy, which was done within 2 weeks, showed longer apnea/hypopnea times in ALS patients who had aspiration than in those without aspiration ( $p = 0.5$ ). But there was no significant difference in apnea/hypopnea times in MD patients with or without aspiration (Fig. 7).

2) Respiratory movement was reduced because of swallowing disturbance: Dysphagia has been reported to aggravate respiratory insufficiency<sup>7) 8)</sup>, because the respiratory muscles are mobilized during swallowing. That could be a reason that apnea/hypopnea elongation occurs during water swallow with severe respiratory failure. Some of the ALS patients and the MD patients who had severely affected respiratory function (lower % FVC) showed longer apnea/hypopnea in our study.

3) Prolongation of oropharyngeal transit<sup>9) 10)</sup>: Oropharyngeal transit time is longer in ALS patients than in MD patients with the same degree of respiratory function (personal study). Long oropharyngeal transit time could lead long respiratory suppression.

Hadjikitis reported that the inhalation phase was likely to occur after swallowing in patients with motor neuron diseases<sup>7)</sup>. It has also been reported that prolonged apnea readily induced inhalation after swallowing<sup>11)</sup>. In another study, a high risk of aspiration during inhalation immediately after swallowing<sup>12)</sup> has been reported. We speculate that prolonged apnea after water swallowing, which leads to inhalation, also increases the risk of aspiration or silent aspiration. In other words, patients who have prolonged apnea are at greater risk of aspiration than those who do not.

This study had some limitations. First, because a thermistor was usually used to measure sleep apnea, the temporal changes from exhalation to inhalation and apnea/hypopnea

were slightly unclear. This limitation precluded evaluation of the occurrence rate of inhalation soon after swallowing. Second, only water was used to test swallowing. The pattern of breathing after a swallow assessed by a water bolus may not be the same as with a chew and swallow, such as semisolid or solid food. Respiratory cycles during and after swallowing have been reported to be irregular depending on the amount of swallowed water or food in the elderly and in patients with chronic obstructive respiratory insufficiency<sup>13)~15)</sup>. Another limitation is that we did not evaluate pneumography and videofluoroscopy simultaneously during water swallowing. However, aspiration does not occur in each swallow. If aspiration induces elongation of apnea/hypopnea, averaged apnea/hypopnea in four swallow times should show the relationship to aspiration.

### Conclusion

Prolonged apnea/hypopnea times were observed during water swallowing in ALS patients. We speculate that this prolongation is caused by severe swallowing disturbance and respiratory failure, which increases the risk of aspiration. The respiration of ALS patients should be closely monitored during eating.

### References

- 1) Subcommittee on Motor Neuron Disease/Amyotrophic Lateral Sclerosis of the World Federation of Neurology Research Group on Neuromuscular Disease, El Escorial "Clinical Limits of Amyotrophic Lateral Sclerosis" Workshop Contributors. El Escorial World Federation of Neurology criteria for the diagnosis of amyotrophic lateral sclerosis. *J Neurol Sci* 1994; 124 (suppl): 96—107
- 2) ALS CNTF Treatment Study (ACTS) Phase I-II Study Group: The amyotrophic lateral sclerosis functional rating scale. *Arch Neurol* 1996; 53: 141—147
- 3) Takahashi K, Groher M, Michi K: Methodology for detecting swallowing sounds. *Dysphagia* 1994; 9: 54—62
- 4) Hirst LJ, Ford GA, Gibson GJ, et al: Swallow-induced al-

- teration in breathing in normal older people. *Dysphagia* 2002; 17: 152—161
- 5) Preiksatis HG, Mills CA: Coordination of breathing and swallowing: effects of bolus consistency and presentation in normal adults. *J Appl Physiol* 1996; 81: 1704—1714
- 6) Nishino T, Yonezawa T, Honda Y: Effects of swallowing on the pattern of continuous respiration in human adults. *Am Rev Respir Dis* 1985; 132: 1219—1222
- 7) Hadjikoutis S, Pickersgill TP, Dawson K, et al: Abnormal patterns of breathing during swallowing in neurological disorders. *Brain* 2000; 123: 1863—1873
- 8) Nozaki S, Kunitomi A, Saito T, et al: Process of swallowing disturbance in amyotrophic lateral sclerosis—Evaluation of videofluoroscopy and respiratory function. *Clin Neurology* 2003; 44: 77—83
- 9) Cha CH, Patten BM: Amyotrophic lateral sclerosis: abnormalities of the tongue on magnetic resonance imaging. *Ann Neurol* 1989; 25: 468—472
- 10) Kawai S, Tsukada M, Mochimatsu I, et al: A study of the early stage of dysphagia in amyotrophic lateral sclerosis. *Dysphagia* 2003; 16: 1—8
- 11) Nilsson H, Ekberg O, Bulow M, et al: Assessment of respiration during video fluoroscopy of dysphagic patients. *Acad Radiol* 1997; 4: 503—507
- 12) Martin BJ, Corlew MM, Wood H, et al: The association of swallowing dysfunction and aspiration pneumonia. *Dysphagia* 1994; 9: 1—6
- 13) Martin BJ, Logemann JA, Shaker R, et al: Coordination between respiration and swallowing: respiratory phase relationship and temporal integration. *J Appl Physiol* 1994; 76: 714—723
- 14) Hiss SG, Treole K, Stuart A: Effect of age, gender, bolus volume, and trial on swallowing apnea duration and swallow/respiratory phase relationships of normal adults. *Dysphagia* 2001; 16: 128—135
- 15) Shaker R, Li Q, Ren J, et al: Coordination of deglutition and phases of respiration. Effect of aging, tachypnea, bolus volume, and chronic obstructive pulmonary disease. *Am J Physiol* 1992; 263: G750—G755